

ANTIPROLIFERATIVE AND ANTIMICROBIAL PROPERTIES OF PURE AND ENCAPSULATED RUTIN

CORINA DANCIU¹, IULIA ANDREEA PINZARU^{2*}, CRISTINA ADRIANA DEHELEAN², MONICA HANCIANU³, ISTVAN ZUPKO⁴, DAN NAVOLAN⁵, MONICA LICKER⁶, ROXANA MARIA GHIULAI⁷, CODRUȚA MARINELA ȘOICA⁷

¹Department of Pharmacognosy, "Victor Babeș" University of Medicine and Pharmacy Timișoara, 2 Eftimie Murgu Square, 300041 Timișoara, Romania

²Department of Toxicology, "Victor Babeș" University of Medicine and Pharmacy Timișoara, 2 Eftimie Murgu Square, 300041 Timișoara, Romania

³Department of Pharmacognosy, University of Medicine and Pharmacy "Gr.T.Popa", Faculty of Pharmacy, 16 University Street, 700115, Iași, Romania

⁴Department of Pharmacodynamics and Biopharmacy, University of Szeged, Eötvös u. 6. Szeged H-6720, Hungary

⁵Department of Obstetrics and Gynaecology, "Victor Babeș" University of Medicine and Pharmacy Timișoara, 2 Eftimie Murgu Square, 300041 Timișoara, Romania

⁶Department of Microbiology-Virology, "Victor Babeș" University of Medicine and Pharmacy Timișoara, 2 Eftimie Murgu Square, 300041, Timișoara, Romania

⁷Department of Pharmaceutical Chemistry, "Victor Babeș" University of Medicine and Pharmacy Timișoara, 2 Eftimie Murgu Square, 300041 Timișoara, Romania

*corresponding author: iuliapinzaru@umft.ro

Manuscript received: September 2017

Abstract

Rutin, a polyphenolic natural compound is described in the literature for a wide range of biological activities. The study aims to test the antiproliferative potential and the antibacterial effect for pure rutin and its β cyclodextrin complexes. On one hand, the antiproliferative assays against A2780 (human ovarian cells), MDA-MB-231 (human breast cancer cells) and SiHa (cervical cancer cells) did not show any substantial inhibition of cell growth. On the other hand, the antimicrobial screening revealed that rutin presents positive effects against: *Streptococcus pyogenes*, *Enterococcus faecalis*, *Bacillus cereus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Escherichia coli*. A decreased, but detectable, activity was noticed for *Staphylococcus aureus* and *Streptococcus pneumoniae*. Incorporation in the ramified cyclodextrins had a mixed effect on the antibacterial activity, with increased, constant or decreased effect, depending on the strain. The study concludes that rutin can be reconsiderate as a natural active antibacterial agent, while modulation of its solubility by incorporation in β -cyclodextrins is debatable depending on the tested strain.

Rezumat

Rutozida este un compus natural cu structură de tip polifenol, descris în literatură pentru o gamă largă de activități biologice. Studiul present și-a propus să testeze potențialul antiproliferativ și efectul antibacterian pentru rutozida pură, respectiv după încorporare în trei tipuri de β ciclodextrine. Pe de o parte testele antiproliferative pe liniile celulare A2780 (celule de cancer ovarian), MDA-MB-231 (celule de cancer de sân) și SiHa (celule de cancer de col uterin) nu au evidențiat nici o inhibare substanțială a creșterii celulare. Pe de altă parte evaluarea antimicrobiană arată că rutozida a prezentat un efect pozitiv împotriva următoarelor tulpini: *Streptococcus pyogenes*, *Enterococcus faecalis*, *Bacillus cereus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* și *Escherichia coli*. O activitate scăzută, dar detectabilă, a fost observată pentru *Staphylococcus aureus* și *Streptococcus pneumoniae*. Incorporarea în ciclodextrine ramificate a avut un efect mixt asupra activității antibacteriene, cu efect crescut, constant sau scăzut, în funcție de tulpină. Studiul concluzionează că rutozida poate fi reconsiderată ca agent antibacterian natural. Modularea solubilității prin încorporarea în ciclodextrinele β ramificate este discutabilă în funcție de tulpina testată.

Keywords: rutin, cyclodextrin, antiproliferative, antimicrobial

Introduction

Rutin is a glycosylated polyphenolic phytochemical which is widespread in coloured fruits and vegetables. Medicinal plants with high rutin content include: buckwheat (*Fagopyrum esculentum* Moench) [17], elderberry (*Sambucus nigra* L.) [13], japanese pagoda

tree (*Sophora japonica* L.) [19] and herb-of-grace (*Ruta graveolens* L.) [22].

The phytochemical was assigned with pleiotropic pharmacological effects (e.g.): capability to increase capillary resistance and decrease capillary permeability [15]; antiplatelet aggregatory effects [9, 10]; important antiinflammatory effects by inhibiting NO, TNF- α

and MPO activity in activated human neutrophils [23] as well the synthesis of pro-inflammatory cytokines, TNF- α and NF- κ B in inflammatory bowel disease [35]. Rutin exhibited powerful antioxidant activity by counteracting the reactive oxygen species (ROS) both *in vitro* [43] and *in vivo* [11] and was demonstrated to have an antiproliferative and proapoptotic activity against different cancer cell lines including: GL-15 human glioblastoma, A549 human lung, MCF-7 breast, HepG2 liver and HT-29 colon cancer cells [33, 42, 44]. Finally, rutin has shown antibacterial and antifungal activity as well as synergistic effects against methicillin resistant *S. aureus* (MRSA), when co-administered with classic antibiotics [1]. Moreover, the phyto-compound was able to reduce the toxicity and improve antifungal activity of Amphotericin B when tested against *Cryptococcus sp.* [25]. Despite the multiple pharmacological effects and its therapeutic potential, rutin exhibits a poor water solubility, thus affecting its *in vivo* bioavailability. For this purpose, many formulation strategies were conducted in order to improve this downside. One of these strategies is represented by encapsulating the flavonoid in cyclodextrins (CDs). CDs are cyclic oligosaccharides made up of α -D-glucopyranoside units, exhibiting a hydrophobic profile on the inside and a hydrophilic one on the outside of the glycosidic ring. Thereby, these features make CDs potential drug delivery systems for hydrophobic molecules in order to increase their bioavailability. Various studies were conducted in order to assess the advantages of CD encapsulation for different types of molecules [12, 36]. In a previous study, several flavonols (fisetin, quercetin and kaempferol) encapsulated in β -CDs showed promising antiproliferative activity *vs.* native compounds when tested on A2780 human ovarian carcinoma, MDA-MB-231 human breast cancer and SiHa human cervix cancer cells [6]. Other types of molecules, such as nerolidol, a component of essential oils, was investigated after encapsulation in α -CD, β -CD, γ -CD, HP- β -CD, RAMEB (randomly methylated-beta-cyclodextrins), CRYSMEB (methylated beta-cyclodextrin) and SBE- β -CD (sulfobutylether- β -cyclodextrin) highlighting their effectiveness as encapsulating agents and solubility enhancers [2]. Also, pentacyclic triterpenes such as betulin esters, ursolic and oleanolic acids complexes with CDs showed increased *in vitro* antiproliferative activity when tested on human melanoma cell lines [26, 30, 31]. All these promising results make this formulation strategy a powerful tool in improving the bioavailability and increasing the effectiveness of hydrophobic phytochemicals with important therapeutic potential.

In the present study, our aim was to investigate native rutin compared to its inclusion complex in CDs regarding: i) the antiproliferative activity on

three tumour cell lines (A2780 human ovarian cancer, MDA-MB-231 human breast cancer and SiHa cervical cancer) and ii) the antibacterial activity against different bacterial strains (*S. pyogenes*, *E. faecalis*, *B. cereus*, *P. aeruginosa*, *K. pneumoniae* and *E. coli*).

Materials and Methods

Reagents

Reagents were purchased as follows: rutin from Sigma Aldrich, Germany, β -cyclodextrin (BCD), hydroxyl-propyl- β -cyclodextrin (HPBCD) and randomly methylated β -cyclodextrin (RAMEB) from Cyclolab, Hungary.

Complexes preparation

For the development of the inclusion complexes the kneading method was used as described in the papers of Danciu *et al.*, [6, 7]. All the complexes were prepared using a molar ration 1:2 active substance: CD ($M_{wRutin} = 610.52$ g/mol, $M_{wBCD} = 1134.98$ g/mol, $M_{wHPBCD} = 1396$ g/mol, $M_{wRAMEB} = 1312$ g/mol).

Cell culture

All three gynaecological cancer cell lines, namely: A2780 (human ovarian), MDA-MB-231 (human breast) and SiHa (cervical) (ECACC-European Collection of Cell Cultures, Salisbury, UK) were cultured in minimal essential medium (MEM), supplemented with 10% foetal calf serum (FCS), 1% antibiotic mixture (Pen/Strep), and 1% nonessential aminoacids (NEAA). All media and supplements were obtained from Lonza Ltd. (Basel, Switzerland). The cells were cultured and expanded in standard conditions, at 37°C and 5% CO₂ atmosphere.

MTT assay

Cells were seeded onto a 96-well culture plate at a cellular density of 5,000 cells/well and attached to the bottom of the well during an overnight. After 24 h, 100 μ L of new medium containing the tested substances (dissolved in dimethyl sulfoxide - DMSO; purchased from Sigma-Aldrich Ltd., Budapest, Hungary) were added and incubated for 72 h. The cells were then assayed by the addition of 20 μ L of 5 mg/mL MTT solution (purchased from Sigma-Aldrich Ltd., Budapest, Hungary). The intact mitochondrial reductase converted MTT into blue formazan crystals during a 4 h contact period and the precipitated crystals were dissolved in 100 μ L DMSO. Finally, the reduced MTT was spectrophotometrically analysed at 545 nm, using a Stat Fax microplate reader (Awareness Technology, Palm City, FL, USA); wells with untreated cells were considered as reference for viability. All *in vitro* experiments were carried out on two microplates in quadruplicates for each substance tested and controls. The DMSO concentration (0.3%) of the medium did not have any significant effect on cell proliferation.

In vitro antibacterial activity

Rutin, its BCD complexes and the native CDs were tested for the antimicrobial activity using 11 strains: *Bacillus cereus* (ATCC 14579), *Staphylococcus aureus* (ATCC 25923), *Streptococcus pyogenes* (ATCC 19615), *Streptococcus pneumoniae* (ATCC 49619), *Enterococcus faecalis* (ATCC 29212), *Escherichia coli* (ATCC 25922), *Yersinia enterocolitica* (ATCC 23715), *Klebsiella pneumoniae* (ATCC 700603), *Proteus mirabilis* (ATCC 25933), *Pseudomonas aeruginosa* (ATCC 27853), *Candida albicans* (ATCC 10231) by the diffusion method as previously described [27, 29]. The antibacterial and antimycotic activity were determinate according to the National Committee for Clinical Laboratory Standards (NCCLS, 1997). A microbial suspension with a concentration of 10^8 CFU/mL was employed. The compounds were diluted in DMSO at a concentration of 10 mM. The inoculum was dispersed on the Mueller-Hinton agar plates and filter paper discs 6 mm in diameter

impregnated with the solution of the tested compounds were distributed on the surface. Paper disks impregnated with BCD, HPBCD and RAMEB were also used. The native CDs did not have any effect on the selected strains. The plates were incubated at 37°C for 24 h. The inhibition zone diameters were determined with a ruler. Triplicate tests were performed and medium values are reported. For positive controls, discs with gentamicin or fluconazole were used, with inhibition zones between 16 - 19 mm.

Results and Discussion

Antiproliferative activity for the selected cell lines A2780, MDA-MB-231 and SiHa showed no substantial growth inhibition when treated with pure rutin and its CDs complexes at 30 µg/mL. In some cases a modest increase in cell viability was evidenced but no practical relevance can be attributed to this phenomenon (Figure 1).




Figure 1.

Antiproliferative activity of rutin and its cyclodextrin inclusion complexes against three tumor cell lines at 30 µg/mL

The antimicrobial screening showed that rutin elicits antibacterial properties on some of the screened strains, namely *S. pyogenes* (19 mm), *E. faecalis* (19 mm), *B. cereus* (16 mm), *P. aeruginosa* (16 mm), *K. pneumoniae* (16 mm) and *E. coli* (16 mm). These strains present comparable inhibition zones with gentamicin, the positive control. A decreased but detectable activity was noticed for *S. aureus* (12 mm) and *S. pneumoniae* (11 mm). Incorporation in the

ramified CDs had a mixt effect on the antibacterial activity. For *S. pyogenes* and *E. faecalis*, incorporation in each of the three CDs conduced to a decreased inhibition zone. For *S. pneumoniae*, *B. cereus*, *P. aeruginosa*, *K. pneumoniae*, and *E. coli* the activity was constant whereas for *S. aureus* the diameter of the inhibition zone increased from 12 mm for pure rutin to 17 mm for rutin:BCD and rutin:HPBCD, respectively 16 mm for rutin: RAMEB (Table I).

Table I

The antimicrobial activity of rutin and its CDs inclusion complexes (the values are presented as average of three determinations)

Strains	Rutin	Rutin BCD	Rutin HPBCD	Rutin RAMEB
<i>E. coli</i>	16 mm	16 mm	16 mm	16 mm
<i>Klebsiella pneumoniae</i>	16 mm	16 mm	16 mm	16 mm
<i>Proteus mirabilis</i>	6 mm	6 mm	6 mm	6 mm
<i>Yersinia enterocolitica</i>	6 mm	6 mm	6 mm	6 mm
<i>Pseudomonas aeruginosa</i>	16 mm	16 mm	16 mm	16 mm
<i>Bacillus cereus</i>	16 mm	16 mm	16 mm	16 mm
<i>Streptococcus pneumoniae</i>	11 mm	11 mm	11 mm	11 mm
<i>Streptococcus pyogenes</i>	19 mm	13 mm	16 mm	15 mm
<i>Staphylococcus aureus</i>	12 mm	17 mm	17 mm	16 mm
<i>Enterococcus faecalis</i>	19 mm	15 mm	11 mm	12 mm

In terms of natural compounds, flavonoids are one of the most studied class of phytochemicals for cancer chemoprevention [7]. Nano-encapsulation represents a method applied in order to enhance bioactivity and/or bioavailability of biologically active molecules. It was demonstrated that internal water molecules of CDs, are displaced by an increased number of hydrophobic compounds [30]. Ghasemzadeh *et al* evaluating the main bioactive compounds from curry leaf extracts (*Murraya koenigii* L.) containing rutin in a concentration of 0.082 mg/g DW as *per* HPLC determination have shown that the extract presents antiproliferative properties against MDA-MB-231 breast cancer cells. The biological effect was assigned to the phytocomplex [10]. It is very well known that rutin is the glycoside of quercetin, and although to the best of our knowledge rutin was not reported in the literature as an active agent on breast, ovarian and cervical cancer cells, quercetin was [32, 38, 40]. Correlated to our results, namely stimulation of proliferation for A2780 cell line, Luo *et al* have described that rutin hydrate stimulates the proliferation of OVCAR-3 ovarian cancer cells [20]. Nevertheless previous studies regarding the *in vitro* and *in vivo* anticancer properties of rutin depict that also this glycoside, not only the aglycone quercetin, acts as an active agent on other cancer cell lines hence our idea to screen its effect on A2780, MDA-MB-231 and SiHa cancer cell lines. Antiproliferative effects were also observed in case of HTC hepatic cells, WEHI-3 murine leukaemia cells, SW480 human colon adenocarcinoma cells and recent papers published informations about its protective effect against induced oxidative stress and apoptosis in human lens epithelial cells as well as against the pro-carcinogenic agent benzo(a)pyrene [5, 18, 46]. Regarding the mechanism of action, rutin acts to various modulators of Wnt signalling. For example, the DNA doxorubicin induced damage was highly diminished when rutin was incubated with HepG2 human liver carcinoma cell line but some studies showed that rutin caused DNA damage on BRCA mutant breast cancer cells [21].

Chemo-preventive effects were detailed in case of animal model of colon cancer and other *in vivo* studies have shown the important effects of rutin in diminishing tumour growth in mice implanted with SW480 human colon adenocarcinoma, metastatic A549 human lung adenocarcinoma, LLC Lewis lung carcinoma and HL-60 human promyelocytic leukaemia [16, 43]. Although active on above mentioned cell lines, rutin showed an overall poor antiproliferative activity against cell lines tested in the present study A2780, MDA-MB-231 and SiHa. Increased dissolution and absorption of rutin may be achieved by complexation with CDs. Paczkowska *et al* have demonstrated that inclusion of rutin in CDs leads to increased chemical solubility, stability, dissolvability and antibacterial activity for some of the selected strains [28]. The antiproliferative effects of rutin proved to be unsubstantial on the selected cell lines which were not changed when it was applied as CD complexes (Figure 1). Some studies have described that the inclusion complex of rutin with HPBCD have been identified to have the highest stabilizing effect in aqueous solutions [24, 37]. The antimicrobial activity of rutin obtained as a methanolic extract from *Pteris vittata* L. was screened against eight intestinal microorganisms by the group of Singh *et al*. Disk diffusion and micro-dilution methods showed efficiency for *P. aeruginosa* and *K. pneumoniae* [34]. Also, aqueous extracts containing rutin from *Castanea sativa* Mill. leaves where analysed in terms of antimicrobial activity against *S. aureus*, *K. pneumoniae*, *P. aeruginosa*, *Proteus vulgaris*, *E. coli*, *Enterobacter cloacae*, *Salmonella typhi* and *Enterobacter aerogens*. Minimum inhibitory concentrations where detected between 32 and 128 µg/mL [3]. In a similar study, Dubey *et al*, assessed the antimicrobial activity of rutin using concentrations ranging from 10 µg/mL to 0.05 µg/mL and reported positive results for *S. aureus*, *E. coli*, *Klebsiella oxytoca*, *P. aeruginosa* and *Candida albicans* [8]. As it can be seen from Table I the results of our screening corroborate with the ones presented by other studies in the field. The novelty that comes with this study is the analyse of

the antimicrobial activity after incorporation in CDs. Wang *et al* have showed an increased antibacterial potency for miconazole after incorporation into a cellulosic fabric grafted with BCD [41]. Sun *et al* have shown that chitosan films containing BCD and essential oils have increased the anti-infective properties of the chitosan films, while Iacovino *et al* have concluded that the antibacterial activity of pipemidic acid is increased for some of the tested strains after incorporation in BCD [14, 39]. In a recent study, Corciova *et al.* proven that complexation of hesperidin with BCD led to higher antibacterial and antioxidant effect [4]. On the other hand, Zhao *et al.* analysing BCD inclusion complex of chlorogenic acid, concluded that in terms of antioxidant and antibacterial capacity no significant difference was observed between the pure substance and the inclusion complex [45]. As depicted in the results section, in this study the majority of the tested strains acted in the same manner after incorporation took place. Only in case of *S. aureus* the potency was increased.

Conclusions

Rutin can be re-considerate as a natural active antibacterial agent against *S. pyogenes*, *E. faecalis*, *B. cereus*, *P. aeruginosa*, *K. pneumoniae* and *E. coli*. Modulation of the solubility by incorporation in CDs is debatable depending on the tested strain. On the other hand, both the glycoside and its CDs inclusion complexes elicited no substantial antiproliferative effect against A2780 human ovarian, MDA-MB-231 human breast and SiHa cervical cancer cell lines at a relatively high final concentration.

Acknowledgement

Authors thank Péter Bérdi for his excellent assistance in the cell based assays.

References

1. Amin MU, Khurram M, Khattak B, Khan J, Antibiotic additive and synergistic action of rutin, morin and quercetin against methicillin resistant *Staphylococcus aureus*. *BMC Complement Altern Med.*, 2015; 15(1): 1-12.
2. Azzi J, Danjou PE, Landy D, Ruellan S, Auezova L, Greige-Gerges H, Fourmentin S, The effect of cyclodextrin complexation on the solubility and photostability of nerolidol as pure compound and as main constituent of cabreuva essential oil. *Beilstein J Org Chem.*, 2017; 13: 835-844.
3. Basile A, Sorbo S, Giordano S, Ricciardi L, Ferrara S, Montesano D, Castaldo Cobianchi R, Vuotto ML, Ferrara L, Antibacterial and allelopathic activity of extract from *Castanea sativa* leaves. *Fitoterapia*, 2000; 71(Suppl.1): S110-S116.
4. Corciova A, Ciobanu C, Poiata A, Mircea C, Nicolescu A, Drobota M, Varganici CD, Pinteala T, Marangoci N, Antibacterial and antioxidant properties of hesperidin:β-cyclodextrin complexes obtained by different techniques. *J Incl Phenom Macrocycl Chem.*, 2015; 81(1-2): 71-84.
5. Cristina Marcarini J, Ferreira Tsuboy MS, Cabral Luiz R, Regina Ribeiro L, Beatriz Hoffmann-Campo C, Sérgio Mantovani M, Investigation of cytotoxic, apoptosis-inducing, genotoxic and protective effects of the flavonoid rutin in HTC hepatic cells. *Exp Toxicol Pathol.*, 2011; 63(5): 459-465.
6. Danciu C, Bojin F, Ambrus R, Muntean D, Soica C, Paunescu V, Cristea M, Pinzaru I, Dehelean C, Physico-chemical and biological evaluation of flavonols: fisetin, quercetin and kaempferol alone and incorporated in beta cyclodextrins. *Anticancer Agents Med Chem.*, 2016; 17(4): 615-626.
7. Danciu C, Citu C, Zupko I, Ghiulai R, Matusz P, Pavel IZ, Popescu A, Pinzaru I, Simu G, Vlad CD, *In vitro* antiproliferative activity of flavonols: Fisetin, quercetin and kaempferol and their cyclodextrin complexes. *Farmacia*, 2015; 63(6): 858-864.
8. Dubey S, Ganeshpurkar A, Bansal D, Dubey N, Experimental studies on bioactive potential of rutin. *Chronicles of Young Scientists*, 2013; 4(2): 153-157.
9. Filip C, Albu E, Lupaşcu D, Filip N, The influence of a new rutin derivative in an experimental model of induced hyperhomocysteinemia in rats. *Farmacia*, 2017; 65(4): 596-599.
10. Ghasemzadeh A, Jaafar HZE., Rahmat A., Devarajan T, Evaluation of bioactive compounds, pharmaceutical quality, and anticancer activity of curry leaf (*Murraya koenigii* L.). *Evid Based Complement Alternat Med.*, 2014; 2014: 1-8.
11. Han X, Xue X, Zhao Y, Li Y, Liu W, Zhang J, Fan S, Rutin-enriched extract from *Coriandrum sativum* L. ameliorates ionizing radiation-induced hematopoietic injury. *Int J Mol Sci.*, 2017; 18(5): 1-20.
12. Heghes A, Hadaruga NG, Fului AV, Bandur GN, Hadaruga DI, Dehelean CA, *Capsicum annuum* extracts/beta-cyclodextrin complexes Thermal analyses-Karl Fischer water titration correlations and antioxidant activity. *J Therm Anal Calorim.*, 2015; 120(1): 603-615.
13. Ho G, Wangenstein H, Barsett H, Elderberry and elderflower extracts, phenolic compounds, and metabolites and their effect on complement, RAW 264.7 macrophages and dendritic cells. *Int J Mol Sci.*, 2017; 18(3): 1-17.
14. Iacovino R, Rapuano F, Caso JV, Russo A, Lavorgna M, Russo C, Isidori M, Russo L, Malgieri G, Isernia C, β-Cyclodextrin Inclusion Complex to Improve Physicochemical Properties of Pipemidic Acid: Characterization and Bioactivity Evaluation. *Int J Mol Sci.*, 2013; 14(7): 13022-13041.
15. Ihme N, Kiesewetter H, Jung F, Hoffmann KH, Birk A, Müller A, Grützner KI, Leg oedema protection from a buckwheat herb tea in patients with chronic venous insufficiency: A single-centre, randomised, double-blind, placebo controlled clinical trial. *Eur J Clin Pharmacol.*, 1996; 50(6): 443-447.
16. Kahali B, Marquez SB, Thompson KW, Yu J, Gramling SJ, Lu L, Aponick A, Reisman D, Flavonoids from each of the six structural groups reactivate BRM, a possible cofactor for the

- anticancer effects of flavonoids. *Carcinogenesis*, 2014; 35(10): 2183-2193.
17. Kreft I, Fabjan N, Yasumoto K, Rutin content in buckwheat (*Fagopyrum esculentum* Moench) food materials and products. *Food Chem.*, 2006; 98(3): 508-512.
 18. Lin JP, Yang JS, Lu CC, Chiang JH, Wu CL, Lin JJ, Lin HL, Yang MD, Liu KC, Chiu TH, Chung JG, Rutin inhibits the proliferation of murine leukemia WEHI-3 cells in vivo and promotes immune response in vivo. *Leuk Res.*, 2009; 33(6): 823-828.
 19. Liu JL, Li LY, He GH, Optimization of microwave-assisted extraction conditions for five major bioactive compounds from flos sophorae immaturus (Cultivars of *Sophora japonica* L.) using response surface methodology. *Molecules*, 2016; 21(3): 1-27.
 20. Luo H, Jiang BH, King SM, Chen YC, Inhibition of cell growth and VEGF expression in ovarian cancer cells by flavonoids. *Nutr Cancer.*, 2008; 60(6): 800-809.
 21. Maeda J, Roybal EJ, Brents CA, Uesaka M, Aizawa Y, Kato TA Natural and glucosyl flavonoids inhibit poly(ADP-ribose) polymerase activity and induce synthetic lethality in BRCA mutant cells. *Oncol Rep.*, 2014; 31(2): 551-556.
 22. Mancuso G, Borgonovo G, Scaglioni L, Bassoli A, Phytochemicals from *Ruta graveolens* activate TAS2R bitter taste receptors and TRP channels involved in gustation and nociception. *Molecules*, 2015; 20(10): 18907-18922.
 23. Nassiri-Asl M, Adineh M, Hajiali F, Nassiri-Asl M, Treatment with rutin - A therapeutic strategy for neutrophil-mediated inflammatory and autoimmune diseases. *J Pharmacopuncture*, 2017; 20(1): 52-56.
 24. Nguyen TA, Liu B, Zhao J, Thomas DS, Hook JM, An investigation into the supramolecular structure, solubility, stability and antioxidant activity of rutin/cyclodextrin inclusion complex. *Food Chem.*, 2013; 136(1): 186-192.
 25. Oliveira VM, Carraro E, Auler ME, Khalil NM, Quercetin and rutin as potential agents antifungal against *Cryptococcus spp.* *Braz J Biol.*, 2016; 76(4): 1029-1034.
 26. Oprean C, Mioc M, Csányi E, Ambrus R, Bojin F, Tatu C, Cristea M, Ivan A, Danciu C, Dehelean C, Paunescu V, Soica C, Improvement of ursolic and oleanolic acids' antitumor activity by complexation with hydrophilic cyclodextrins. *Biomed Pharmacother.*, 2016; 83: 1095-1104.
 27. Oprean C, Zambori C, Borcan F, Soica C, Zupko I, Minorics R, Bojin F, Ambrus R, Muntean D, Danciu C, Pinzaru IA, Dehelean C, Paunescu V, Tanasie G, Anti-proliferative and antibacterial in vitro evaluation of the polyurethane nanostructures incorporating pentacyclic triterpenes. *Pharm Biol.*, 2016; 54(11): 2714-2722
 28. Paczkowska M, Mizera M, Piotrowska H, Szymanowska-Powalowska D, Lewandowska K, Goscianska J, Pietrzak R, Bednarski W, Majka Z, Cielecka-Piontek J, Complex of rutin with β -cyclodextrin as potential delivery system. *PLoS One* 2015, 10(3): 1-16.
 29. Pavel IZ, Danciu C, Oprean C, Dehelean CA, Muntean D, Csuk R, Muntean DM, *In vitro* evaluation of the antimicrobial ability and cytotoxicity on two melanoma cell lines of a benzylamide derivative of maslinic acid. *Analytical Cellular Pathology*, 2016; 2016: 1-6.
 30. Pinzaru I, Hadaruga DI, Hadaruga NG, Corpas L, Grozescu I, Peter F, Hepatoprotective flavonoid bioconjugate/beta-cyclodextrin nanoparticles: DSC - molecular modeling correlation. *Dig J Nanomater Biostruct.*, 2011; 6(4): 1605-1617.
 31. Pinzaru I, Trandafirescu C, Szabadai Z, Mioc M, Ledeti I, Coricovac D, Ciurlea S, Ghiulai RM, Crainiceanu Z, Simu G, Synthesis and biological evaluation of some pentacyclic lupane triterpenoid esters. *Rev Chim.*, 2014; 65(7): 848-851.
 32. Ren MX, Deng XH, Ai F, Yuan GY, Song HY, Effect of quercetin on the proliferation of the human ovarian cancer cell line SKOV-3 in vitro. *Exp Ther Med.*, 2015; 10(2): 579-583.
 33. Santos BL, Silva AR, Pitanga BP, Sousa CS, Grangeiro MS, Fragomeni BO Coelho PL, Oliveira MN, Menezes-Filho NJ, Costa MF, El-Bachá RS, Velozo ES, Sampaio GP, Freire SM, Tardy M, Costa SL, Antiproliferative, proapoptotic and morphogenic effects of the flavonoid rutin on human glioblastoma cells. *Food Chem.*, 2011; 127(2): 404-411.
 34. Singh M, Govindarajan R, Rawat AKS, Khare PB, Antimicrobial flavonoid rutin from *Pteris vittata* L. against pathogenic gastrointestinal microflora. *American Fern Journal.*, 2008; 98(2): 98-103.
 35. Singh UP, Singh NP, Busbee B, Guan H, Singh B, Price RL, Taub DD, Mishra MK, Nagarkatti M, Nagarkatti PS, Alternative medicines as emerging therapies for inflammatory bowel diseases. *Int Rev Immunol.*, 2012; 31(1): 66-84.
 36. Soica C, Coricovac D, Dehelean C1, Pinzaru I, Mioc M, Danciu C, Fulas A, Puiu M, Sitaru C, Nanocarriers as tools in delivering active compounds for immune system related pathologies. *Rec Pat Nanotechnol.*, 2016; 10(2): 128-145.
 37. Sri KV, Kondaiah A, Ratna JV, Annappurna A, Preparation and characterization of quercetin and rutin cyclodextrin inclusion complexes. *Drug Dev Ind Pharm.*, 2007; 33(3): 245-253.
 38. Srivastava S, Somasagara RR, Hegde M, Nishana M, Tadi SK, Srivastava M, Choudhary B, Raghavan SC, Quercetin, a natural flavonoid interacts with dna, arrests cell cycle and causes tumor regression by activating mitochondrial pathway of apoptosis. *Sci Rep.*, 2016; 6: 1-13.
 39. Sun X, Sui S, Ference C, Zhang Y, Sun S, Zhou N, Zhu W, Zhou K, Antimicrobial and mechanical properties of β -cyclodextrin inclusion with essential oils in chitosan films. *J Agric Food Chem.*, 2014; 62(35): 8914-8918.
 40. Vidya Priyadarsini R, Senthil Murugan R, Maitreyi S, Ramalingam K, Karunakaran D, Nagini S, The flavonoid quercetin induces cell cycle arrest and mitochondria-mediated apoptosis in human cervical cancer (HeLa) cells through p53 induction and NF- κ B inhibition. *Eur J Pharmacol.*, 2010; 649(1-3): 84-91.
 41. Wang J, Cai Z, Incorporation of the antibacterial agent, miconazole nitrate into a cellulosic fabric grafted with β -cyclodextrin. *Carbohydr Polym.*, 2008; 72(4): 695-700.

-
42. Xu DP, Zheng J, Zhou Y, Li Y, Li S, Li HB, Extraction of natural antioxidants from the *Thelephora ganbajun* mushroom by an ultrasound-assisted extraction technique and evaluation of antiproliferative activity of the extract against human cancer cells. *Int J Mol Sci.*, 2016; 17(10):1-15.
43. Yang J, Guo J, Yuan J, *In vitro* antioxidant properties of rutin. *LWT - Food Sci Technol.*, 2008; 41(6): 1060-1066.
44. Yang K, Lamprecht SA, Liu Y, Shinozaki H, Fan K, Leung D, Newmark H, Steele VE, Kelloff GJ, Lipkin M, Chemoprevention studies of the flavonoids quercetin and rutin in normal and azoxymethane-treated mouse colon. *Carcinogenesis*, 2000; 21(9): 1655-1660.
45. Zhao M, Wang H, Yang B, Tao H, Identification of cyclodextrin inclusion complex of chlorogenic acid and its antimicrobial activity. *Food Chem.*, 2010; 120(4): 1138-1142.
46. Zhou YF, Guo B, Ye MJ, Liao RF, Li SL, Protective effect of rutin against H₂O₂-induced oxidative stress and apoptosis in human lens epithelial cells. *Curr Eye Res.*, 2016; 41(7): 933-942.